

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau


6869

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/00, 15/64, 15/67 C12N 15/70, 15/54	A1	(11) International Publication Number: WO 92/14819 (43) International Publication Date: 3 September 1992 (03.09.92)
(21) International Application Number: PCT/US92/01074 (22) International Filing Date: 21 February 1992 (21.02.92) (30) Priority data: 662,224 26 February 1991 (26.02.91) US (71) Applicant: E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors: PIERCE, James, Clifford ; 2644 Bellows Drive, Wilmington, DE 19810 (US). STERNBERG, Nat, Leon ; 1116 Dorset Drive, West Chester, PA 19380 (US). (74) Agents: GALLEGOS, R., Thomas et al.; E.I. du Pont de Nemours and Company, Legal/Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		(81) Designated States: AT (European patent), BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), MC (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: A POSITIVE SELECTION VECTOR FOR THE BACTERIOPHAGE P1 CLONING SYSTEM**(57) Abstract**

Positive selection cassettes are disclosed which contain a lethal gene, a promoter, a repressor sequence overlapping the promoter, and a cloning site between the promoter and the lethal gene. Insertion of a foreign nucleic acid sequence into the cloning site prevents expression of the lethal gene. Expression of the lethal gene under nonrepressed conditions kills a host organism containing a positive selection cassette which does not contain the foreign nucleic acid sequence.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

TITLEA POSITIVE SELECTION VECTOR FOR THE
BACTERIOPHAGE P1 CLONING SYSTEM

5

FIELD OF THE INVENTION

This invention allows the selection of large molecular weight DNA inserts in the P1 cloning system by positive selection for the ability of clones with inserts to grow on media in the presence of sucrose.

10 This new P1 vector also facilitates characterization of cloned DNA.

BACKGROUND OF THE INVENTION

The bacteriophage P1 cloning system allows the headful in vitro packaging of foreign DNA fragments as large as 95 kb in length. Sternberg, Proc. Natl. Acad. Sci. USA 87, 103-107 (1990) has shown that the P1 cloning system can generate 100,000 clones containing inserts per microgram of vector DNA. Large molecular weight clones are faithfully replicated in E. coli host strains and DNA from these clones can be easily isolated by standard molecular biological techniques. Thus, the P1 cloning system rivals Yeast Artificial Chromosomes (YAC) and cosmid cloning systems for the generation and characterization of genomic libraries.

25 Cosmid cloning vectors were designed by Bruning et al., Gene 4, 85-107 (1978) and Collins et al., Proc. Natl. Acad. Sci. USA, 75, 4242-4246 (1978), so that a bacteriophage lambda in vitro packaging reaction can encapsulate insert DNA up to 47 kb and infect E. coli at high efficiency. The cosmid vector plus insert DNA is cyclized in the E. coli bacterium at the lambda cos sites located on the vector. The same cos site is used in recognition by the lambda packaging apparatus for encapsulation of the vector-insert DNA into the lambda bacteriophage head. A major limitation of the cosmid

30

35

cloning system is the relatively small size of the insert clone (47 kb). Many eukaryotic genes have been shown to be larger than 50 kb with some genes (e.g., dystrophin) up to 1000 kb. The small size of cosmid clones necessitates a labor intensive and "error-prone" procedure of multiple chromosome "walking" and "jumping" methodologies when isolating large genomic clones.

Another system for cloning large molecular weight DNA fragments is Yeast Artificial Chromosomes (YAC's) developed by Burke et al., Science 236, 806-812 (1987). YAC cloning enables DNA inserts up to 1000 kb to be propagated as minichromosomes in specific yeast strains. YAC vectors contain a yeast replication origin, a centromere, and a set of telomeres. After ligation of insert DNA to the YAC vector, the DNA is introduced into yeast spheroplasts by direct DNA transformation. The major limitation of YAC cloning is the inefficiency of the transformation reaction (about 1000 clones per microgram of vector DNA) and the difficulty in characterization of YAC clones once they have been generated. The YAC clone represents a small proportion (less than 1%) of the total DNA in a yeast cell. This makes recovery, isolation, and analysis of any particular YAC clone burdensome.

The bacteriophage P1 cloning system complements both cosmid and YAC cloning in the construction of genomic libraries. A 50,000-member human DNA library has been generated in the P1 cloning system by Sternberg et al., The New Biol. 2, 151-162 (1990) which represents about a one times coverage of the human genome. The most recent P1 cloning vector (pNS582tet14Ad10) consists of a P1 pac site used for the initiation of headful packaging, two P1 lox sites which cyclize the P1 vector upon introduction in an *E. coli* host strain containing the P1 cre protein, a kanamycin gene for determining

which *E. coli* cells contain a P1 plasmid, and a tetracycline gene for the cloning of insert DNA. The P1 cloning vector also contains a bacteriophage P1 plasmid replicon which maintains the P1 clone at a single copy per cell, and an IPTG inducible P1 lytic replicon for amplifying P1 clones in DNA isolation procedures. Another aspect of the cloning vector is a 10 kb "stuffer fragment" from adenovirus DNA which gives flexibility in the headful packaging reaction.

10 A model P1 cloning reaction consists of cutting the pNS582tet14Ad10 with the restriction enzymes ScaI and BamHI to generate 5 kb and 25 kb vector "arms". The digested vector DNA is then treated with calf intestine alkaline phosphatase to inhibit self ligation of the
15 vector. The vector arms are added to genomic DNA fragments that were previously digested with a BamHI-end compatible restriction enzyme (e.g. Sau3A). The two DNA's are then ligated and a portion of the ligation mixture is added to the first part of the two stage P1
20 in vitro packaging reaction. The first reaction consists of a cell extract prepared from P1 infected *E. coli* which is enriched for the P1 pac cleavage proteins. After pac cleavage, the DNA mixture is incubated in the stage II P1 in vitro packaging reaction which consists
25 of a *E. coli* cell extract enriched for P1 virion capsids and tails. The phage encapsulated DNA is then infected into an *E. coli* host strain that contains the cre recombinase. A lox-lox site specific recombination reaction effectively cyclizes the P1 vector-insert clone
30 which is maintained as a single copy extrachromosomal circular plasmid. To isolate DNA from a P1 clone, the cell containing the clone is grown in the presence of IPTG which induces the P1 lytic replicon. This induction increases the copy number of the P1 clone about
35 25 fold which gives enough DNA (about 1 microgram) from

a 10 ml mini-alkaline lysis DNA isolation procedure for standard restriction mapping and size characterization procedures.

One problem encountered in the P1 cloning system is that a significant number of P1 vector molecules that contained no insert were present after a typical cloning experiment. These "no-insert" clones interfered with subsequent analysis of the cloning experiment in two ways. First, the number of clones to be screened when looking for a particular DNA insert was markedly increased due to the presence of "no-insert" containing clones. Secondly, upon subsequent growth of *E. coli* from a P1 cloning experiment, the bacteria that contained a "no-insert" vector generally grew much better than clones that contained large DNA inserts. Therefore, after a few rounds of growth the population of *E. coli* containing clones was greatly increased for "no-insert" vector clones.

To overcome the problems encountered in the previous versions of the P1 cloning system the pAd10-SacBII positive selection P1 cloning vector was developed. Many other positive selection based cloning systems have been developed for standard plasmid based recombinant DNA work. Henrich et. al., Gene 42, 345-349 (1986) demonstrated a positive selection vector based on the E gene (lysis protein) of bacteriophage ϕ X174. Kuhn et. al., Gene 42, 253-263 (1986) developed a system which uses the EcoRI endonuclease. Burns et. al., Gene 27, 323-325 (1984) showed that positive selection can be generated in a system based on resistance to 5-fluorouracil. Other similar systems are listed in the Burns et al. article.

Another positive selection system used in DNA cloning is based on the sacB gene from Bacillus subtilis. This gene codes for the enzyme levansucrase

(sucrose:2,6- β -D-fructan 6- β -D fructosyltransferase; EC 2.4.1.10) which catalyzes the transfructorylation of sucrose to various acceptor substrates resulting in the hydrolysis of sucrose and levan synthesis. Gay et al., J. Bacteriol. 164, 918-921 (1985) demonstrated that the production of levansucrase in *E. coli* is lethal in the presence of growth media containing 5% sucrose. Gay et al. have used this knowledge to develop a positive selection cloning system based on inactivating the *B. subtilis* sacB structural gene. This allows the growth of only those *E. coli* bacteria containing recombinant clones that have DNA inserts when grown in the presence of sucrose.

Tang et. al., Gene (in press) (1990), (U.S. Patent Application Serial No. 07/376,474) have cloned the sacB gene from *Bacillus amyloliquefaciens* and shown extensive DNA sequence homology to the sacB gene from *B. subtilis*. When the sacB gene from *B. amyloliquefaciens* was cloned on a multicopy plasmid in *E. coli*, a lethal phenotype is observed when cells are grown in the presence of sucrose. This knowledge has inspired us to develop a novel P1 positive selection cloning vector, pAd10-SacBII.

SUMMARY OF THE INVENTION

This invention describes a new bacteriophage P1 cloning vector (pAd10-SacBII) which allows for the positive selection of clones containing large molecular weight inserts. Improvements of the P1 cloning system are:

- (a) a positive selection system based on the sacB gene from *Bacillus amyloliquefaciens* which codes for the enzyme levansucrase (sucrose:2,6- β -D-fructan 6- β -D fructosyltransferase; E.C.2.4.1.10). A unique promoter cassette has been invented which

allows the disruption of sacB expression by cloning foreign DNA into a unique BamHI site between the promoter and sacB structural gene.

- 5 (b) a regulatory system based on the bacteriophage P1 c1 repressor protein to control the expression of the lethal sacB gene.
- 10 (c) an E. coli host strain that contains an integrated lambda phage that expresses the P1 c1 gene.
- 15 (d) placement of a bacteriophage T7 and Sp6 RNA polymerase promoters which border the cloning site of the P1 vector to facilitate characterization and analysis of P1 clones.
- 20 (e) placement of unique rare cutting restriction enzyme sites, SfiI SalI, and NotI which border the BamHI cloning site of the P1 vector to facilitate characterization and isolation of cloned foreign DNA inserts.

BRIEF DESCRIPTION OF THE FIGURES

25 Figure 1 illustrates the steps involved in constructing the P1 positive selection vector.

Figure 2 illustrates the positive selection P1 cloning vector pAd10-SacBII.

30 Figure 3 illustrates a BglII/XhoI restriction digest of pAd10-SacBII-human clones from (sucrose) and (no sucrose) agar plates.

Figure 4 illustrates the ability to linearize and size characterize a P1 clone using the rare restriction site NotI.

STATEMENT OF DEPOSIT

The following plasmids and bacteria relating to this invention have been deposited in the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852-1776 under the Budapest Treaty.

pAd10-SacBII was designated ATCC Accession No. 68505.

pAd10-SacBI was designated ATCC Accession No. 68504.

10 NS3607 was designated ATCC Accession No. 55135.

DETAILED DESCRIPTION OF THE INVENTION

The vector described herein is an improvement to the P1 cloning system in the following manner:

(1) The pAd10-SacBII vector allows for the positive selection of large molecular weight DNA clones by killing clones that do not have a DNA insert when bacteria containing the clone are grown on media supplemented with sucrose.

(2) Analysis of large molecular weight clones is facilitated by the ability to remove the vector DNA fragment away from the clone insert by cutting with the restriction enzymes NotI and SfiI or SalI which are unique to the vector and directly border the unique BamHI site used in cloning.

(3) Chromosomal "walking" and "jumping" procedures will be greatly improved by the ability to make RNA and DNA probes from both ends of the DNA insert by using the bacteriophage T7 and Sp6 promoters that border the unique BamHI site.

(4) DNA sequence information from both ends of the cloned DNA insert will be readily obtained by using the unique bacteriophage T7 and Sp6 promoters as sites for DNA sequencing primers. This sequence information can then be reported as Sequence Tagged Sites (STS) for each P1 clone.

Construction of the pAd10-SacBII Vector

Construction of the P1 positive selection cloning vector (pAd10-SacBII) was initiated by cutting the parent P1 vector pNS582tet14Ad10 (Figure 1), The New Biol. 2, 151-162 (1990), available from New England Nuclear as NENPHAGE®, with the restriction enzymes SalI and BamHI. This removed a 276 base pair fragment from the tetracycline gene of the parent vector. A synthetic duplex DNA oligonucleotide (promoter cassette) Lewin, B. (1983) Genes. John Wiley and Sons, NY, was then inserted into the SalI-BamHI site in a two step process. First, a 52 base pair duplex oligonucleotide. (The upper strand is referred to as SEQ ID NO. 1; the lower strand is referred to as SEQ ID NO. 2.)

5' TCGAGCTTGA CATTGTAGGA CTATATTGCT CTAATAAATT TCGGCGCGCT TG 3'
CG AACTGTAACA TCCTGATATA ACGAGATTAT TTAAACGCCG GCGAACCTAG

that contained a consensus *E. coli* promoter sequence, a P1 c1 repressor sequence that overlapped the *E. coli* synthetic promoter, and a unique NotI site, was inserted into the SalI and BamHI site of the parent vector while regenerating the BamHI site but destroying the SalI site. Next, a 37 base pair duplex synthetic oligonucleotide. (The upper strand is referred to as SEQ ID NO. 3; the lower strand is referred to as SEQ ID NO. 4.)

5' GGCCGCGGAT CCGTCGACGG CCAATTAGGC CTACGTA 3'
CGCCTAG GCAGCTGCCG GTTAATCCGG ATGCATCTAG

that contained a series of restriction sites (NotI, BamHI, SalI, SfiI, SnaBI), was inserted into the first 52 base pair oligonucleotide using a NotI and BamHI digest. The NotI site was regenerated while the BamHI

site was destroyed, leaving only a unique BamHI site located between the NotI and SalI sites. This series of genetic engineering steps created a P1 plasmid construct named pAd10-37 as shown in Figure 1.

5 The next step in the construction of the P1 positive selection vector was to transfer a 5.0 kilobase (kb) ClaI DNA fragment which includes the pAd10-37 tetracycline gene with the promoter cassette insert, P1 lytic replicon, and part of the kanamycin gene into the
10 ClaI site of pBR322. This was done to facilitate further experimental manipulations with a smaller plasmid (9 kb) rather than the 30 kb pAd10-37 plasmid. This plasmid was named pBR5.0. The pAd10-37 plasmid and the plasmid pBR322 were cut with the ClaI restriction
15 enzyme. The 5.0 kilobase fragment was isolated and ligated into the ClaI site of pBR322.

 A pUC plasmid (pBE501) (Figure 1) containing the sacB gene from Bacillus amyloliquefaciens (gift from Vansantha Nagarajan, E. I. du Pont de Nemours and
20 Company, CR&D Microbiology, U.S. Patent Application Serial No. 07/376,474) was cut with restriction enzyme EcoRI which generated a 1.6 kb DNA fragment that contains the structural gene for sacB and the ribosome binding site but does not contain the endogenous sacB
25 promoter. A 9 base pair EcoRI/SnaBI adapter. (The upper strand is referred to as SEQ ID NO. 5; the lower strand is referred to as SEQ ID NO. 6.)

30 5' CCACTAGTC 3'
 GGTGATCAGTTAA

 that contains an internal unique SpeI restriction site was placed on each end of the 1.6 kb EcoRI DNA fragment. This was done so that the sacB structural gene was
35 bordered by unique SpeI sites for easy physical

identification and characterization of clones containing this insert. The modified 1.6 kb sacB fragment was ligated via blunt ends, into the unique SnaBI site of pBR5.0. DNA from this ligation reaction was transformed
5 into E. coli strain DH5 α Iq (available from Bethesda Research Labs) that already contained a plasmid pACYC-P1cl, New England Biolabs. The pACYC-P1cl plasmid contains the gene for the P1 cl repressor protein, which is needed to prevent expression of the sacB gene, which
10 even under permissive growth conditions (no sucrose in the media) exhibits a lethal phenotype. The P1 cl protein is expressed by bacteriophage P1 to repress phage lytic functions during vegetative growth. This protein acts by binding an asymmetric recognition DNA
15 sequence, usually in vicinity of an RNA polymerase promoter, Eliason and Sternberg, J. Mol. Biol. 198, 181-293 (1987). The lethal phenotype is probably due to the strong consensus E. coli promoter regulating the over expression of the SacB gene to produce a periplasmic
20 SacB protein. The P1 cl repressor protein is able to block expression of the sacB gene and allow replication of the otherwise lethal plasmid. Positive clones were identified by hybridization against the nick translated, radioactively labeled 1.6 kilobase fragment from plasmid
25 pBE501. The correct plasmid construct was confirmed by physical mapping with diagnostic restriction enzymes and by the sensitivity of E. coli DH5 α Iq cells containing the plasmid to growth media containing sucrose, even in the presence of the P1 cl repressor protein. This
30 plasmid construct was named pBR5.0-17.

The next step in the construction of the P1 positive selection vector was to return the 6.6 kb ClaI fragment (previously 5.0 kb) of plasmid pBR5.0-17 to the parent pAd10-37 vector. DNA from plasmid pBR5.0-17 was
35 subjected to a partial ClaI restriction digest, since

there is a ClaI site in the sacB structural gene, and
th 6.6 kb fragment was isolated and ligated to the 25
kb ClaI fragment of pAd10-37. DNA from this ligation
reaction was transformed into E. coli strain DH5 α Iq that
5 contained a lambda prophage expressing the P1 c1 gene.
Positive clones were identified by restoration of the
kanamycin resistant phenotype and by physical mapping
via restriction enzyme digests. This P1 vector was
named pAd10-SacBI (Figure 1). Initial characterization
10 of bacterial cells containing this vector showed a
lethal phenotype when grown on media containing sucrose.
When DNA inserts were cloned into the unique BamHI site
of pAd10-SacBI, cells containing these constructs were
imparted with the ability to grow on media containing
15 sucrose. This result demonstrated the utility and
success of the P1 SacB positive selection vector.

To further increase the utility of the P1 SacB
positive selection vector, a modification of the unique
restriction site region (NotI, BamHI, SalI, and SfiI)
20 was performed. The pAd10-SacBI vector was cut with the
restriction enzymes NotI and SalI and a 56 base pair
duplex synthetic oligonucleotide containing the promoter
sequences for the T7 RNA polymerase and Sp6 RNA
polymerase with a new BamHI site between the promoters
25 was inserted into the NotI/SalI site at a position which
directly bordered the unique BamHI cloning site, thus
deleting the original unique BamHI site. (The upper
strand is referred to as SEQ ID NO. 4; the lower strand
is referred to as SEQ ID NO. 8.)

30

```
5'  GGCCGCTAAT ACGACTCACT ATAGGGAGAG GATCCTTCTA TAGTGTCACC TAAATG   3'  
    CGATTA TGCTGAGTGA TATCCCTCTC CTAGGAAGAT ATCACAGTGG ATTTACAGCT
```

The resulting fragment is referred to in SacBII (Figure
35 1). The plasmid containing SacBII is referred to as

pAd10-SacBII (Figure 2). Confirmation of the correct plasmid construct was generated by restriction mapping and testing for the functional presence of the T7 and Sp6 promoters. We also observed that cells containing the pAd10-SacBII vector exhibited a "less" lethal phenotype when grown without the P1 cI repressor in the absence of sucrose. This may be due to an "attenuator-like" effect produced by the insertion of the T7 and Sp6 promoter DNA sequences between the sacB structural gene and its synthetic promoter. This is the final version of the P1 positive selection vector and is named pAd10-SacBII. Experiments describing the utility of this vector are described below.

Construction of E. coli host strain
expressing the P1 cI repressor

To replicate the pAd10-SacBII vector the expression of levansucrase must be controlled. sacB gene expression under the control of the synthetic E. coli promoter gives a lethal phenotype in the absence of sucrose. We therefore constructed a host strain which expresses the P1 cI gene at a high enough level to inhibit levansucrase expression. This was accomplished by constructing a bacteriophage lambda prophage that contains the P1 cI repressor gene and inserting the construct into E. coli DP501q.

Phage lambda (imm21-P1:7A5b) is a phage containing a functioning P1 cI repressor gene. It was generated from the starting phage lambda (dam15 b575 b529 attλ+ imm21) in which P1 EcoRI fragment 7 (O'Brien, Genetic Maps, Cold Spring Harbor Press, Cold Spring Harbor, NY 1990) was cloned at lambda map coordinate 65.8. EcoRI-7 contains the P1 cre gene, the P1 loxP site, and the p1 cI repressor gene, Sternberg et al., J. Mol. Biol. 187, 197-212 (1986). Deletion mutants of the above starting phage were isolated and the location of the deleted DNA

determined by restriction mapping, as described in Eliason and Sternberg, J. Mol. Biol. 198, 281-293 (1987). One mutant, designated lambda (imm21-P1:7Δ5b), contained a contiguous deletion of DNA extending from the attλ site into the cloned P1 sequences. P1 BamHI-EcoRI fragment 8* (cre containing EcoRI-BamHI fragment from P1 EcoRI fragment 7) was completely removed and part of P1 BamHI fragment 9 was removed. In vivo loxP-cre reactions indicated that this lambda deletion mutant construct was loxP-, cre-. Further experiments showed that an *E. coli* lysogen containing this prophage synthesized functional P1 c1 repressor protein. This strain is called NS3607. These results indicate that the Δ5b deletion terminates between P1 loxP and the P1 c1 gene in the P1 BamHI fragment 9 (O'Brien, supra). The *E. coli* DH5αIq (P1 c1) was used as a host strain to prepare pAd10-SacBII plasmid DNA for cloning experiments described below.

Preparation of pAd10-SacBII DNA

Significant care must be given in the preparation of pAd10-SacBII DNA. Vector DNA is prepared by the cesium chloride density gradient method as described by Godson and Vapnek, Biochim. Biophys. Acta. 299, 516-522 (1973). The observation that bacteria containing the SacBI vector expressed a lethal phenotype even in the absence of sucrose in the media resulted in a careful analysis of cesium chloride prepared SacBI DNA. Restriction analysis showed that a significant portion of the vector DNA was deleted in the region containing the sacB gene. When this DNA was transformed into *E. coli* DH5αIq and the bacteria grown on agar plates containing 5% sucrose, only bacteria that contained the population of SacBI plasmids that had a deletion grew well. This result significantly compromises the utility

of the P1 positive selection system since it generated a large background of clones not containing DNA inserts, exactly what the SacB positive selection was supposed to eliminate. This problem was overcome by the

5 construction of the pAd10-SacBII vector which contains an extra 43 base pairs of DNA (T7 and Sp6 RNA promoters) between the sacB structural gene and its synthetic promoter. E. coli DH5 α Iq (P1 cl) NS3607 that contained this vector (SacBII) did not display the lethal

10 phenotype in the absence of sucrose in the growth media. Still, careful analysis of vector DNA is necessary when preparing large amounts of DNA for cloning experiments. This is achieved by restriction analysis of vector DNA and by transformation of competent E. coli cells with

15 vector DNA to determine the percentage of the plasmid population which are sucrose resistant.

Restriction digest of pAd10-SacBII DNA

Vector arms are prepared by a sequential double restriction digest of the SacBII DNA. Three micrograms

20 of vector DNA was incubated with the restriction enzyme ScaI (New England Biolabs) as per manufacture's instructions, in a 30 microliter reaction. The reaction was incubated at 37°C for one hour. The reaction volume was increased to 50 microliters with water, BamHI

25 restriction buffer and 1 microliter of BamHI. The reaction was incubated for 10 minutes at 37°C. The digest reaction was then extracted with one volume phenol, one volume chloroform-isoamyl alcohol (24:1) saving the aqueous layer each time. The DNA^a was

30 precipitated with two to three volumes of ice cold ethanol and 0.3 M sodium acetate and the mixture placed on ice for 30 minutes. The DNA was collected by centrifugation in a microcentrifuge at room temperature for 20 minutes, washed with 0.5 ml of 75% ethanol and

35 the pellet air dried. The DNA pellet was resuspended in

20 microliters of TE buffer (10 mM Tris (pH 8.0) 1 mM EDTA).

Preparation and analysis of high molecular weight insert DNA for cloning into pAd10-SacBII

5 There are many protocols for the preparation of genomic DNA that can be used in the P1 cloning system. This depends upon the organism, tissue or cell type in question. The pAd10-SacBII vector was tested on DNA prepared from human lymphoblastoid cell line 697 as
10 described by Sternberg, U.S. Patent Application No. 07/397,071, (1989) and Sternberg et. al., The New Biol. 2, 151-162 (1990). Briefly, DNA was isolated from lysed cells and fractionated on a 10-40% sucrose gradient. After dialysis, 0.5 ml of the large molecular weight DNA
15 was incubated with 4 units of Sau3I restrictive enzyme and restriction buffer minus magnesium, overnight at 4°C. This ensured adequate mixing of the Sau3I restriction enzyme with the viscous genomic DNA. Digestion was initiated by adding magnesium chloride to
20 10 mM and aliquots were removed at 10 different time points and the reaction terminated by adding EDTA to 20 mM and heating at 70°C for 15 minutes.

 The Sau3I partial digest of genomic DNA was analyzed by removing 10% of each time point fraction and
25 subjecting the DNA to agarose field inversion pulse gel electrophoresis. The DNA was fractionated via electrophoresis to resolve fragment sizes between 20 and 200 kb by a 1% agarose gel in 0.5 X's TBE buffer (89 mM Tris pH 8.0, 89 mM borate, 2 mM ethylene diamine
30 tetraacetic acid (EDTA)), for 4 hours at 180 volts with a 0.6 sec. forward, 0.2 sec. reverse, and ramp of 20, pulse conditions. The gel was stained with ethidium bromide and DNA visualized with uv fluorescence. Those aliquots that contained digested genomic DNA fragments

in the 70 t 100 kb rang were us d as substrat for cloning into the pAd10-SacBII vector.

Ligation and Packaging of SacBII vector
and genomic insert DNA

5 The ligation of the SacBII vector arms to genomic DNA and its subsequent in vitro packaging into P1 capsids has been described in detail in Sternberg, Proc. Natl. Acad. Sci. USA 87, 103-107 (1990) and Sternberg, U.S. Patent Application No. 07/397,071. 0.2 micrograms
10 ScaI/BamHI digest SacBII vector arms were incubated with Sau3A digested genomic DNA or with TE buffer (no insert control experiment) in ligase buffer (50 mM Tris pH 7.5, 10 mM MgCl₂, 10 mM dithiothreitol, 50 mg/ml bovine serum albumin) for 30 minutes at 37°C. DNA ligase and ATP (to
15 2 mM) were then added and the reaction incubated at 16°C overnight. The ligation reaction was then heated at 65°C for 5 minutes and added to the first part of the two stage in vitro P1 packaging reaction. The stage 1 extract is a bacteriophage P1 lysate that contains the
20 enzymes needed to cleave the pac site of the P1 cloning vector. The stage 2 extract is a bacteriophage P1 lysate that contains the components (phage capsids and tails) necessary to form an infectious phage particle. The stage 1 reaction was incubated for 15 minutes at
25 30°C and then transfered to the stage 2 reaction, which was incubated for 20 minutes at 30°C. The final reaction was diluted to a total volume of 180 microliters with TMG buffer (10 mM Tris pH 8.0, 10 mM MgCl₂, 0.1% gelatin) plus pancreatic DNase (10 ug/ ml).
30 Fifteen microliters of chloroform were added to the P1 packaging reaction, and then stored at 4°C.

Growth and DNA preparation of P1 clones
containing high molecular weight DNA inserts

A 10 to 20 microliter aliquot from each P1
35 packaging reaction was added to 0.1 ml of mid-log phase,

concentrated (10X), *E. coli* bacteria containing the cre recombinase (strains NS3145 or NS3529). After a 10 minute incubation at 37°C, one ml of L broth was added to the phage/cell mixture and then incubated for 30-60 minutes at 37°C with shaking. The infected cells were pelleted in a microcentrifuge and then resuspended in 0.15 ml of L broth and then spread on L agar plates that contained 25 ug/ml kanamycin with or without 5% sucrose. The plates were incubated at 37°C overnight. Kanamycin and kanamycin/sucrose resistant colonies were recorded.

P1 clone plasmid DNA was prepared by picking a single bacterial colony with a 50 microliter capillary pipet and adding it to 10 ml of L broth containing 25 ug/ml kanamycin. The cells were grown at 37°C for about 3 hours (early log phase) and IPTG was added to a final concentration of 1 mM. The cells were grown for another 5 hours and then pelleted. DNA was prepared by the alkaline lysis procedure of Birnboim et al., Nucl. Acids Res. 7, 1513-1523 (1979). Plasmid DNA was resuspended in 40 microliters of TE buffer and 0.4 micrograms RNase.

Characterization of genomic insert DNA
from pAd10-SacBII P1 clones

The pAd10-SacBII vector allows the characterization of insert DNA by use of the novel rare cutting restriction sites which border the BamHI cloning site and by the ability to make RNA probes from either or both ends of the genomic insert via the T7 and Sp6 promoters. Physical mapping and size characterization of cloned insert DNA is facilitated by the ability to isolate the SacBII vector sequence away from the genomic DNA. Chromosome walking and DNA sequencing of cloned inserts is achieved by the utilization of the unique promoter sequences. Those skilled in the art will recognize that these features have been successfully

utilized in plasmid and cosmid cloning as described in Melton et al., Nucl. Acids Res. 12, 7035-7056 (1984) and Wahl et al., Proc. Natl. Acad. Sci. USA 84, 2160-2164 (1987).

5 Physical characterization of P1 clones was carried out by restriction enzyme digestion. Each digestion reaction contains 13 microliters of plasmid DNA from a particular p1 clone, 1.5 microliters of restriction buffer, and 1.0 microliter of each restriction enzyme
10 (either BglII, NotI or NotI and SalI). The restriction digest was performed at 37°C for 2 hours, then subjected to a phenol/chloroform extraction, and air dried to remove excess chloroform. If the volume was too large an ethanol precipitation step was also included and the
15 DNA resuspended in 15 microliters of TE buffer. The DNA was then analyzed on a 1% agarose pulse field inversion gel electrophoresis in 0.5X TBE buffer using pulse conditions of 0.6 sec. forward, 0.2 sec. reverse, ramp 20 at 120 volts for 15 hours. The gel was then stained
20 with ethidium bromide and the DNA visualized by uv fluorescence.

EXAMPLE 1

Transformation efficiencies of uncut and cut/ligated pAd10-SacBII

25 To determine the efficiency of the positive selection feature of the pAd10-SacBII vector, DNA transformation experiments were carried out. After a cesium chloride banded DNA preparation of the SacBII vector had been generated, the DNA was analyzed by a
30 restriction digest with BglII, SpeI, and BamHI/ScaI. The results showed that the correct size DNA fragments were produced with no apparent rearrangements observed. Two hundred nanograms of pAd10-SacBII from two different DNA preparations was transformed into *E. coli* strains
35 NS3145 (cre+) and DH5Iq (cre-) as described by Maniatis

et al., Molecular cloning: A Laboratory Manual (Cold Spring Harbor Laboratory, Box 100, Cold Spring Harbor, NY (1982)). Strain NS3145 maintains the SacBII vector as a single copy plasmid due to cre mediated lox-lox recombination. Strain DH5Iq maintains the SacBII vector as a multicopy plasmid. This difference is important since the sacB positive selection function must work with only one copy of the gene present per bacterial cell. The transformed cells were plated on L agar plates containing kanamycin or kanamycin/5% sucrose and grown overnight at 37°C. The number of resistant colonies were counted (or estimated) and are shown in part A of Table 1. The same pAd10-SacBII DNA (100 ng each) was cut with the restriction enzymes BamHI, SalI, SacI/BamHI or ScaI/SalI, extracted with phenol/chloroform, ethanol precipitated and resuspended in TE buffer (as described previously). The digested DNA was then placed in a ligation reaction, incubated overnight at 16°C, and transformed into E. coli strains NS3145 and DH5Iq (as described above). Kanamycin and kanamycin/sucrose resistant colonies were recorded and are shown in part B of Table 1.

TABLE 1
DNA Transformation Efficiencies of pAd10-SacBII

5	DNA	NS3145 (cre+)		DH5Iq (cre-)	
		Kan ^r	Sucrose ^r	Kan ^r	Sucrose ^r
	A)				
	uncut #1	~10,000	6	~10,000	2
	uncut #2	~10,000	7	~10,000	7
10	B)				
	uncut	361	2	405	0
	BamHI	135	6	90	5
	SalI	246	5	222	3
	ScaI/BamHI	56	7	9	13
15	SacI/SalI	107	7	34	9

The results in part (A) show that the positive selection efficiency, calculated as the number of kan-r colonies divided by the number of kan/sucrose-r colonies, for the uncut SacBII vector was approximately 1000 fold. This was true whether the sacB gene was at single (NS3145) or multicopy (DH5Iq). These results indicate that only a small proportion of the pAd10-SacBII DNA contains an inactive sacB gene or nonfunctional synthetic promoter. This "background" population is most likely due to single base mutations in the sacB structural gene and promoter region which change the expression of levansucrase either quantitatively or qualitatively, thus allowing growth on media containing sucrose. The results from part (B) indicate that molecular biological manipulations (restriction digestion and ligation reactions and phenol/chloroform extractions) do have an impact on the efficiency of the SacB positive selection system. The transformation efficiency was decreased even for the uncut DNA, although less DNA (100 ng as opposed to 200

ng) was used for each strain transformed. The positive selection efficiency was approximately 200 to >400 fold. This number dropped somewhat when the pAd10-SacBII was cut at the unique BamHI and Sall sites between the sacB structural gene and the synthetic promoter. Since a ligation step is necessary to achieve a functional pAd10-SacBII molecule, it is possible that plasmid dimers can form which inactivate the sacB gene by a head to head or tail to tail arrangement. These plasmid dimers would still be kan-r but would no longer show sucrose sensitivity. When the SacBII vector was cut to generate "arms" via the ScaI/BamHI or Sall double digest, transformation efficiencies were markedly decreased, this is not surprising since ScaI creates a blunt end which is recalcitrant to ligation. Interestingly the positive selection efficiency was very poor (2 to 5 fold). This may be due to the fact that under these conditions most of the kan-r colonies are from generated plasmids via aberrant ligation products that inactivate the sacB gene hence giving a sucrose resistant phenotype.

EXAMPLE 2

DNA Packaging and Positive Selection

Efficiencies of pAd10-SacBII

To determine how well the positive selection system of the SacBII vector would perform during the construction of a genomic library, the following model experiment was done. Four micrograms of pAd10-SacBII DNA was cut with the appropriate restriction enzyme (as described above) at 37°C for 3 hours. DNA was extracted with phenol/chloroform and then ethanol precipitated. Some samples were then treated with calf intestinal phosphatase (CIP) (see Table 2, part (A) below) by resuspending the DNA in 50 microliters of 50 mM Tris (pH 8.0), 0.1 mM EDTA. Calf intestinal phosphatase was

added to a final concentration of 0.01 units and the reaction incubated at 37°C for 1 hour. The CIP reaction was extracted with phenol/chloroform, ethanol precipitated and resuspended in 20 microliters of TE buffer. The cut SacBII vector DNA was then placed in a ligation reaction in the absence of foreign DNA, under the following conditions: ~3 ug vector DNA (18 ul), 3 ul 10X ligase buffer, 2 ul 25 mM ATP, 1.5 ul T4 DNA ligase, 5 ul TE buffer. The DNA was first heated at 70°C for 2 minutes and then the rest of the ligation reaction added and incubated at 16°C overnight. The ligation reaction was then heated at 70°C for 3 minutes and added to the two stage P1 in vitro packaging reaction as described previously.

15 In part (B) of Table 2, partially Sau3A digested human DNA was used in the ligation reaction with the pAd10-SacBII vector. Two hundred nanograms of vector DNA was digested with ScaI and BamHI and added to Sau3A partially digested human genomic DNA (as described previously) and incubated at 37°C for 30 minutes. Ligation reactions were set up as follows: 1.5 ul (200 ng) vector DNA, 15 ul (~1 ug) human genomic DNA, 2 ul 10X ligase buffer, 1 ul 25 mM ATP, 1 ul T4 DNA ligase and incubated overnight at 16°C. Tris-EDTA buffer was substituted for the vector control reaction. 25 Three different time fractions (6, 8, and 10 minutes) of the human genomic Sau3A partial digest were used. The ligation reactions were heated for 3 minutes at 70°C and added to the P1 in vitro packaging reaction as described previously. The number of kanamycin and kanamycin/sucrose resistant colonies were recorded and are presented in Table 2.

TABLE 2
DNA Packaging and Positive Selection
Efficiency of pAd10-SacBII

5	Vector DNA	Insert DNA	NS3529 (cre+) Kan ^r	Sucrose ^r
	A)			
	uncut	none	~2000	1
10	BamHI	none	~1500	200
	ScaI	none	~2000	0
	BamHI (CIP)	none	~250	5
	ScaI (CIP) /BamHI	none	~1500	42
	ScaI/BamHI (CIP)	none	~350	28
15	B)			
	ScaI/BamHI	none	800	8
	ScaI/BamHI	Hu-6"	686	516
	ScaI/BamHI	Hu-8"	518	328
	ScaI/BamHI	Hu-10"	574	418
20				

These results show that the SacBII vector significantly decreases the amount of kan-r colonies when plated on media containing 5% sucrose. In part (A) of Table 2 the SacBII vector was cut with restriction enzymes BamHI or ScaI and then religated. Interestingly, only the BamHI cut increased the number of sucrose resistant colonies while the ScaI cut, which is distant from the sacB structural gene, did not. This data is in accord with the data from Table 1 which showed an increase in the number of sucrose resistant colonies when the region between the sacB gene and its promoter were manipulated via restriction digest and ligation. Phosphatase treatment with CIP did decrease the total number of kanamycin resistant colonies when the BamHI site was treated. Also, CIP treatment

improved the ratio of sucrose-r colonies to kan-r colonies.

Part (B) of Table 2 reflects the experimental conditions in which genomic libraries can be generated using the P1 cloning system. As stated previously, one problem with the old P1 cloning vector (pAd10) was the number of kan-r colonies generated with no insert. As can be seen from the control ScaI/BamHI with no insert DNA the number of kan-r colonies are decrease 100 fold when plated on media containing sucrose. This result demonstrates that the SacBII vector greatly reduces the contribution of potential "no-insert" containing clones during a genomic cloning experiment. This result is further supported by the data generated when genomic DNA is added to cut SacBII vector during the ligation reactions. The number of sucrose-r colonies is increased over 60 fold in the Hu-6" reaction. Also, the relative number of sucrose-r colonies divided by kan-r colonies is greatly decreased indicating that most of the SacBII vector has productively ligated to genomic DNA fragments. It is that fraction of "insert-less" clones which grow on the kan-r plate but not on the sucrose-r plate that the SacBII positive cloning system was designed to select against. The following example demonstrates that the sucrose-r clones do contain genomic DNA while some of the kan-r clones do not.

EXAMPLE 3

Analysis and Characterization of SacBII

High Molecular Weight Human Clones

To demonstrate that the pAd10-SacBII sucrose resistant colonies actually contain genomic DNA inserts, DNA was prepared from a number of colonies from both the kan-r and sucrose-r populations generated in the packaging experiment described in Example 2. These clones were analyzed via restriction mapping with

BglII/XhoI and fractionation on agarose gel electrophoresis. The SacBII-human DNA's were also characterized by NotI restriction digest and fractionation via pulse field inversion gel electrophoresis.

DNA was prepared (as described above) from 64 colonies from the kanamycin/5% sucrose plates and 12 colonies from the kanamycin plates from the Hu-6" and Hu-8" reactions described in Example 2. Thirteen microliters of DNA was digested with BglII and XhoI at 37°C for 1 hour. The reaction was stopped by heating at 70°C for 5 minutes in the presence of stop dye buffer (6X = 0.25% bromophenol blue, 0.25% xylene cyanol, 40% (w/v) sucrose in water) and the DNA fragments were then fractionated on a 1% agarose gel for 12 hours at 20 volts in 1X TBE buffer. The gels were stained with ethidium bromide and DNA visualized by uv fluorescence. Results show that almost all (>95%) of the sucrose-r clones contain a genomic DNA insert but that many of the kan-r clones (~66%) do not contain DNA inserts. This difference is due to the positive selection aspect of the pAd10-SacBII vector.

A photograph of a gel that is a composite of a representative portion of the sampled colonies is presented in Figure 3. Lane 1 contains molecular weight markers, lanes 2-11 contain DNA from colonies derived from the kanamycin/sucrose agar plates, lanes 12-15 contain DNA from colonies from the kanamycin plates. The DNA from clones in lanes 2-4 are SacBII vectors that contain relatively small genomic inserts. Lanes 5-11 are SacBII clones that contain large molecular weight human genomic inserts. Lanes 12-15 contain DNA with no genomic DNA inserts and reflect the DNA fragments generated from the pAd10-SacBII vector after it has gone through cre mediated lox-lox recombination. Background

fragments are due to the presence of the F' plasmid that contains the *lacIq* gene recombinase in *E. coli* strain NS3145.

To further characterize the SacBII P1 clones DNA from a portion of the clones presented in Figure 3 were subjected to a NotI restriction digest and then fractionated on a 1% agarose field inversion gel electrophoresis (methodology as described above). A photograph of this gel is presented in Figure 4. Lanes 1, 15-18 are molecular weight markers, lanes 2-5 are clones from kanamycin plates (no sucrose), lanes 6-14 are clones from the kanamycin/sucrose agar plates. The clones from the no sucrose plates contain no genomic DNA inserts and the size of the SacBII fragment after lox-lox recombination is about 18 kb. The clones from the plus sucrose plates contain DNA inserts of two general sizes. The large molecular weight clones range in size from 75 to 100 kb. The low molecular weight clones are in the 30 to 50 kb range. DNA's in lanes 2-8 are from strain NS3145 and those in lanes 9-14 are from strain NS3529. The smaller NotI fragment in lane 8 is from the F'*lacIq* plasmid in strain NS3145 and the larger fragment is from the SacBII-human plasmid. These results demonstrate the power of rare restriction site mapping (in this case NotI) in the analysis of P1 clones using the new pAd10-SacBII vector.

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention and without departing from the spirit and scope thereof, can make various modifications and changes of the invention to adapt to various uses and conditions.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- i) APPLICANT: Nat L. Sternberg, Ph.D.
James C. Pierce, Ph.D.
- (ii) TITLE OF INVENTION: A Positive Selection Vector
for the Bacteriophage P1
Cloning System
- (iii) NUMBER OF SEQUENCES: 8
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: E. I. du Pont de Nemours and
Company
 - (B) STREET: 1007 Market Street
 - (C) CITY: Wilmington
 - (D) STATE: Delaware
 - (E) COUNTRY: U.S.A.
 - (F) ZIP: 19898
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette, 3.50 inch, 1.0MB
 - (B) COMPUTER: Macintosh
 - (C) OPERATING SYSTEM: Macintosh System Software,
Version 6.0
 - (D) SOFTWARE: Microsoft Word, Version 4.0
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: UNKNOWN
 - (B) FILING DATE: UNKNOWN
 - (C) CLASSIFICATION: UNKNOWN
- (vii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: R. Thomas Gallegos
 - (B) REGISTRATION NUMBER: 32,692
 - (C) Reference/Docket Number: CR-8930
- (viii) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (302) 892-7342
 - (B) TELEFAX: (302) 892-7949
 - (C) TELEX: 835420

SEQ LISTING

(2) INFORMATION FOR SEQ ID NO:1:

- (i) Sequence Characteristics:
 - (A) LENGTH: 52 bases
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: synthetic oligonucleotide
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (v) ORIGINAL SOURCE:
 - (A) ORGANISM: not applicable
 - (B) STRAIN: not applicable
 - (C) CELL TYPE: not applicable
- (vi) IMMEDIATE SOURCE:
 - (A) LIBRARY: not applicable
 - (B) CLONE: not applicable
- (vii) FEATURES, SEQUENCE IDENTIFICATION SEQ ID NO:1:
- (viii) PUBLICATION INFORMATION:
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TCGAGCTTGA CATTGTAGGA CTATATTGCT CTAATAAATT TCGGGCCGCT TG

(2) INFORMATION FOR SEQ ID NO:2:

- (i) Sequence Characteristics:
 - (A) LENGTH: 52 bases
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: synthetic oligonucleotide
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (v) ORIGINAL SOURCE:
 - (A) ORGANISM: not applicable
 - (B) STRAIN: not applicable
 - (C) CELL TYPE: not applicable

- (vi) IMMEDIATE SOURCE:
 - (A) LIBRARY: not applicable
 - (B) CLONE: not applicable

(vii) FEATURES, SEQUENCE IDENTIFICATION SEQ ID NO:2:

(viii) PUBLICATION INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GATCCAAGCG GCCGCAATT TATTAGAGCA ATATAGTCCT ACAATGTCAA GC

(2) INFORMATION FOR SEQ ID NO:3:

- (i) Sequence Characteristics:
 - (A) LENGTH: 37 bases
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: synthetic oligonucleotide
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (v) ORIGINAL SOURCE:
 - (A) ORGANISM: not applicable
 - (B) STRAIN: not applicable
 - (C) CELL TYPE: not applicable
- (vi) IMMEDIATE SOURCE:
 - (A) LIBRARY: not applicable
 - (B) CLONE: not applicable

(vii) FEATURES, SEQUENCE IDENTIFICATION SEQ ID NO:3:

(viii) PUBLICATION INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGCCGCGGAT CCGTCGACGG CCAATTAGGC CTACGTA

(2) INFORMATION FOR SEQ ID NO:4:

- (i) Sequence Characteristics:
 - (A) LENGTH: 37 bases
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: synthetic oligonucleotide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) ORIGINAL SOURCE:

- (A) ORGANISM: not applicable
- (B) STRAIN: not applicable
- (C) CELL TYPE: not applicable

(vi) IMMEDIATE SOURCE:

- (A) LIBRARY: not applicable
- (B) CLONE: not applicable

(vii) FEATURES, SEQUENCE IDENTIFICATION SEQ ID NO:4:

(viii) PUBLICATION INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GATCTACGTA GGCCTAATTG GCCGTCGACG GATCCGC

(2) INFORMATION FOR SEQ ID NO:5:

(i) Sequence Characteristics:

- (A) LENGTH: 9 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic oligonucleotide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) ORIGINAL SOURCE:

- (A) ORGANISM: not applicable
- (B) STRAIN: not applicable
- (C) CELL TYPE: not applicable

(vi) IMMEDIATE SOURCE:

- (A) LIBRARY: not applicable
- (B) CLONE: not applicable

(vii) FEATURES, SEQUENCE IDENTIFICATION SEQ ID NO:5:

(viii) PUBLICATION INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CCACTAGTC

(2) INFORMATION FOR SEQ ID NO:6:

- (i) Sequence Characteristics:
 - (A) LENGTH: 13 bases
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: synthetic oligonucleotide
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (v) ORIGINAL SOURCE:
 - (A) ORGANISM: not applicable
 - (B) STRAIN: not applicable
 - (C) CELL TYPE: not applicable
- (vi) IMMEDIATE SOURCE:
 - (A) LIBRARY: not applicable
 - (B) CLONE: not applicable
- (vii) FEATURES, SEQUENCE IDENTIFICATION SEQ ID NO:6:
- (viii) PUBLICATION INFORMATION:
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

AATTGACTAG TGG

(2) INFORMATION FOR SEQ ID NO:7:

- (i) Sequence Characteristics:
 - (A) LENGTH: 56 bases
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: synthetic oligonucleotide
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (v) ORIGINAL SOURCE:
 - (A) ORGANISM: not applicable
 - (B) STRAIN: not applicable
 - (C) CELL TYPE: not applicable
- (vi) IMMEDIATE SOURCE:
 - (A) LIBRARY: not applicable
 - (B) CLONE: not applicable

(vii) FEATURES, SEQUENCE IDENTIFICATION SEQ ID NO:7:

(viii) PUBLICATION INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GGCCGCTAAT ACGACTCACT ATAGGGAGAG GATCCTTCTA TAGTGTCCACC TAAATG

(2) INFORMATION FOR SEQ ID NO:8:

(i) Sequence Characteristics:

(A) LENGTH: 56 bases

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic oligonucleotide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) ORIGINAL SOURCE:

(A) ORGANISM: not applicable

(B) STRAIN: not applicable

(C) CELL TYPE: not applicable

(vi) IMMEDIATE SOURCE:

(A) LIBRARY: not applicable

(B) CLONE: not applicable

(vii) FEATURES, SEQUENCE IDENTIFICATION SEQ ID NO:8:

(viii) PUBLICATION INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TCGACATTTA GGTGACACTA TAGAAGGATC CTCTCCCTAT AGTGAGTCGT ATTAGC

What is claimed is:

1. A positive selection cassette, comprising:
 - 5 (a) a gene, the expression of which is lethal to a host cell when the host cell is grown under nonrepressed conditions;
 - (b) a synthetic consensus Escherichia coli promoter upstream from the lethal gene;
 - 10 (c) a P1 c1 repressor sequence overlapping the promoter; and
 - (d) a cloning site between the lethal gene and the promoter, the cloning site bordered by an Sp6 and a T7 RNA promoter, wherein
15 cloning a nucleic acid into the cloning site prevents expression of the lethal gene.
2. A positive selection cassette, comprising:
 - 20 (a) a SacB gene from Bacillus amyloliquefaciens;
 - (b) a synthetic consensus Escherichia coli promoter upstream from the SacB gene;
 - 25 (c) a P1 c1 repressor sequence overlapping the promoter; and
 - (d) a cloning cassette comprising Sp6 and T7 promoters, bordered by BamHI restriction site, the Sp6 and T7 promoters, SfiI rare restriction site and a NotI restriction
30 site, the cloning site located between the synthetic consensus promoter and the SacB gene, wherein cloning of a nucleic acid at the cloning site prevents expression of the SacB gene allowing growth of the host
35 organism in a medium containing sucrose.

3. A vector comprising the positive selection cassette of Claim 1.

5 4. A positive selection cassette, comprising:
 (a) a gene, the expression of which is lethal
 to a host cell when the host cell is grown
 under nonrepressed conditions;
 (b) a synthetic consensus Escherichia coli
10 promoter upstream from the lethal gene;
 (c) a P1 cl repressor sequence overlapping the
 promoter; and
 (d) a cloning site between the lethal gene and
 the promoter, wherein cloning a nucleic
15 acid into the cloning site prevents
 expression of the lethal gene.

 5. A vector comprising the positive selection
 cassette of Claim 4.
20

 6. A host cell genetically engineered to comprise
 a P1 cl repressor sequence.

 7. A host cell comprising a P1 cl repressor
25 sequence and a positive selection cassette.

 8. The host cell of Claim 7 further comprising a
 vector which comprises the P1 cl repressor sequence.

30 9. The host cell of Claim 7 further comprising a
 vector which comprises the positive selection cassette.

 10. The host cell of Claim 7 wherein the positive
 selection cassette is the SacBI cassette.
35

11. The host cell of Claim 7 where in the positive selection cassette is the SacBII cassette.

12. A method of positive selection of nucleic acid clones, comprising:


- (a) cloning a large molecular weight DNA fragment into the vector of Claim 3 at the cloning site;
- (b) inserting the vector into an appropriate host cell; and
- (c) growing the host cell under nonrepressed conditions.

13. A method of positive selection of nucleic acid clones, comprising:

- (a) cloning a large molecular weight DNA fragment into the vector of Claim 5 at the cloning site;
- (b) inserting the vector into an appropriate host cell; and
- (c) growing the host cell under nonrepressed conditions.



1/5

FIG. 1A



(CONTINUED FROM FIG. 1A) 2/5


FIG. 1B




SUBSTITUTE SHEET

3/5

FIG. 2




4/5
FIG.3



SUBSTITUTE SHEET

FIG. 4




SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 92/01074

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C12N15/00; C12N15/64; C12N15/67; C12N15/70 C12N15/54		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C12N	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	JOURNAL OF BACTERIOLOGY vol. 164, no. 2, November 1985, AM. SOC. MICROBIOL., BALTIMORE, US; pages 918 - 921; P. GAY ET AL.: 'Positive selection procedure for entrapment of insertion sequence elements in gram-negative bacteria' cited in the application ---	1-13
A	GENE vol. 55, 1987, ELSEVIER PUBLISHERS, N.Y., U.S.; pages 67 - 74; P.E. STEVIS AND N.W.Y. HO: 'Positive selection vectors based on xylose utilization suppression' --- -/-	1-13
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
10 JUNE 1992	24. 06. 92	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	HORNIG H. 	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	GENE vol. 42, 1986, ELSEVIER PUBLISHERS, N.Y., U.S.; pages 345 - 349; B. HENRICH AND R. PLAPP: 'Use of lysis gene of bacteriophage phiX174 for the construction of a positive selection vector' cited in the application ---	1-13
A	GENE vol. 44, 1986, ELSEVIER PUBLISHERS, N.Y., U.S.; pages 253 - 263; I. KUHN ET AL.: 'Positive selection vectors utilizing lethality of the EcoRI endonuclease' cited in the application ---	1-13